

Pollution Status Of Chandlodia Lake Located In Ahmedabad-Gujarat

Pradeep U. Verma*, Anshita R. Purohit **and Naimesh J. Patel***

(Assistant Professor, Shri R. R. Lalan College – Bhuj-Kachchh, Kachchh University, Gujarat (India) - 370001
(Senior Research Fellow, Genetic Diagnostic center, Department of Zoology, Gujarat University, Ahmedabad, Gujarat (India) – 380009

(C.E.O., Bhishma Agri Research Biotech Co., Biotech Department, Ahmedabad, Gujarat (India) -380009

ABSTRACT

Pollution is viewed as the release of substances and energy as waste product of human activities which result in harmful changes within the natural environment. In the present study water of Chandlodia lake was analyzed for various physico-chemical parameters. The study was carried out for a period of one year. Monthly data's been collected and were represented seasonally along with standard error. Different parameters taken in the study were Temperature, Electrical conductivity, Turbidity, Total dissolve solids, pH, Alkalinity, Total Hardness, Calcium, Magnesium, Dissolved Oxygen, Biochemical oxygen demand, Chloride, Sodium, Nitrate and Phosphate. Therefore from the above result it was concluded that water of Chandlodia lake shows very high level of pollution.

1. INTRODUCTION

Pollutants are defined as the substances which cause pollution. They can be physical or chemical. Each lake system is unique, and its dynamics can be understood only to a limited degree based on information from other lakes. Just as a physician would not diagnose an individual's medical condition or prescribe treatment without a personal medical examination, a limnologist or hydrologist cannot accurately assess a lake system or suggest a management strategy without data and analysis from that particular lake and its environment. The study was carried out to check the pollution status of Chandlodia lake. Chandlodia lake is located in Ahmedabad.

Chandlodia lake is an artificial lake, constructed by AUDA. Chandlodia lake is also located in the western part of Ahmedabad city. The half portions of the lake become dry during the summer season. The people in the surrounding region use to throw waste into the lake and the cattle also take bath in the lake. The lake covers an area of 22,940 m². And its exact geographical location is 23°05'00.05" N Latitude and 72°33'09.43" E Longitude.

2. METHODOLOGY

The present study was carried out for Chandlodia Lake, located in Ahmedabad city. In the

present study the sampling was done during morning hour. The water samples were collected in the polyethylene bottles. The closed bottle was dipped in the lake at the depth of 0.5 to 0.7 m, and then a bottle was opened inside and was closed again to bring it out at the surface. The samples were collected from five different points and were mixed together to prepare an integrated sample. From the time of sample collection to the time of actual analyses, many physical and chemical reactions would change the quality of the water sample; therefore to minimize this change the sample were preserved soon after the collection. The water samples were preserved by adding chemical preservatives and by lowering the temperature. The water temperature, pH, DO, EC and TDS were analyzed immediately on the spot after the collection, whereas the analyses of remaining parameters were done in the laboratory.

The study was carried for a period of 1 year (March 2010 to February 2011). Monthly data was collected, but results were represented season wise. Four month make one season [March to June summer season, July to October monsoon season, and November to February winter season]. The collected water samples were brought to the laboratory and relevant analysis was performed. pH was determined electrometrically using digital pH meter, electrical conductivity was measured by conductivity meter, dissolved oxygen is measured by DO meter, total dissolve solid was measured by using TDS meter and similarly turbidity is measured by Nephthalo turbidity meter. Alkalinity, chloride, TDS, calcium, magnesium, total hardness, nitrate and phosphate were determined by method suggested by [1] [2] [3]. Estimation of sodium was done by Flame Photometric method. The mean value of the monthly data was calculated as season wise and standard error was also calculated by using following formula

Standard deviation

$$\sigma = \sqrt{\frac{\sum(x_i - m)^2}{n-1}}$$

Standard error

$$\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$$

3. RESULT AND DISCUSSION

3.1 Temperature

The temperature plays a crucial role in physico-chemical and biological behavior of aquatic system [4]. Temperature is one of the most important factors in the aquatic environment [5]. The temperature of Chandlodia lake ranges between 16 ± 1.47 to 29 ± 1.58 . The maximum temperature was recorded during summer season and minimum was recorded during winter season.

Generally water temperature correspond with air temperature indicating that the samples collected from shallow zone has a direct relevance with air temperature, shallow water reacts quickly with changes in atmospheric temperature [6] [7] [8] [9] [10] [11].

3.2 Electrical Conductivity

Electrical conductivity in the water is due to salt present in water and current produced by them. It measures the electric current which is proportional to mineral matter present in water. A high level of conductivity reflects on the pollution status as well as trophic levels of the aquatic body [12]. Conductivity of water depends upon the concentration of ions and its nutrient status and variation in dissolve solid content. Electrical conductivity recorded in Chandlodia lake ranges between 2.78 ± 0.07 to 3.24 ± 0.26 . The high value of conductivity was recorded during monsoon season were as low value was recorded during summer season.

The water during the summer decrease as a result some of the aquatic plant got destroyed and very few plants remain in the water. Therefore aquatic plants are in very rare amount in monsoon season, thus electrical conductivity is more in monsoon because water is free from vegetation and aquatic life therefore all the ion are accumulated in water. The decomposition of plants and animals released the ion back in water [13].

3.3 Turbidity

Turbidity is the measure of the light scattered by suspended particles. It is due to the substances not present in the form of solution. Clay, slit, organic matter, phytoplankton and other microscopic organisms cause turbidity in pond water [14]. Light penetration is also highly affected by turbidity. Turbidity in Chandlodia lake recorded ranges between 16 ± 1.47 to 22 ± 0.91 . The maximum turbidity in water was recorded during monsoon season and minimum turbidity was recorded during summer season.

High turbidity in lake water during monsoon season is due to addition of sand, clay, slit, dung and various other pollutant along with rain water from the surrounding area into the lake [15] [16] [17]. High turbidity during monsoon season may be due to inflow of storm water from the surrounding area [18] [19].

3.4 Total Dissolve solids

Total dissolved solids denote mainly the various kinds of mineral present in the water. Dissolved solid do not contain any gas and colloids. In natural water dissolved solids are composed mainly of carbonates, bicarbonates of calcium, magnesium, sodium, potassium, iron and manganese etc. The amount of total dissolve solid in Chandlodia lake ranges between 963 ± 31.5 to 1264 ± 39 . The maximum amount of total dissolve solid was recorded during monsoon season and minimum was recorded during winter.

The high value of TDS during monsoon may be due to addition of domestic waste water, garbage and sewage etc in the natural surface water body. Indeed, high concentration of TDS enriches the nutrient status of water body which was resulted into eutrophication of aquatic ecosystem [5] [20].

3.5 pH

pH measure the concentration of hydrogen ion in water. It is the measurement of acidity or alkalinity. Generally in India many small confined water pockets particularly, are alkaline in nature [21]. The pH value ranges between 8.7 ± 0.17 to 9.1 ± 0.18 . The maximum pH was recorded during monsoon and minimum pH was recorded during summer season.

High pH values promote the growth of algae and results in heavy bloom of phytoplankton [22] [23] [24] [25]. The pH values above 8 in natural water are produced by photosynthetic rate that demand more CO_2 than quantities furnished by respiration and decomposition [26]. The pH of water also depends on the relative quantities of calcium, carbonate and bicarbonate.

3.6 Alkalinity

Alkalinity in natural water is due to free hydroxyl ion and hydrolysis of salts formed by weak acid and strong bases and also due to salt containing carbonates and bicarbonates silicate and phosphate along with hydroxyl ion in the Free states. The change in alkalinity depend on carbonates and bicarbonates, which in term depend upon release of CO_2 . Change in carbonates and bicarbonates also depend upon release of CO_2 through respiration of living organisms. The amount of total alkalinity in Chandlodia lake ranges between 192 ± 3.65 to 214 ± 6.87 . The minimum value of alkalinity was recorded during summer season and maximum value was recorded during monsoon season

The addition of large amount of sewage waste and organic pollutant in the lake also effect photosynthesis rate, which also result in death of plants and living organism. The degradation of plants, living organism and organic waste might also be one of the reasons for increase in a carbonate and bicarbonate, resulting an increase in alkalinity

value [27] [28] [29]. Increase in alkalinity during monsoon was due to input of freshwater and dissolution of calcium carbonate ions in the water column [30].

3.7 Total Hardness

Water hardness is the traditional measure of the capacity of water to react with soap, hard water requiring a considerable amount of soap to produce lather. Hardness of water is not a specific constituent but is a variable and complex mixture of cations and anions. The total hardness recorded in the water of Chandlodia lake ranges between 304 ± 6.58 to 328 ± 8.29 . The maximum amount of total hardness in the water of Chandlodia lake was recorded during monsoon season and minimum amount was recorded during winter season.

Total hardness is mainly due to calcium magnesium and eutrophication [31]. The high value of hardness during monsoon may be due to presence of high content of calcium and magnesium in addition to sulphate and nitrate in the sewage waste added during monsoon [32] [33].

3.8 Calcium

Calcium is an important nutrient for aquatic, organism and it is commonly present in all water bodies [16]. The amount of calcium in the water of Chandlodia lake ranges between 66 ± 3.92 to 82 ± 4.32 . The maximum amount of calcium in the water of Chandlodia lake was recorded during monsoon season and minimum amount was recorded during winter season.

Calcium is present in water naturally, but the addition of sewage waste might also be responsible for the increase in amount of calcium [32] [34]. The decrease in amount of calcium may be due to its absorption by living organisms.

3.9 Magnesium

Magnesium is found in various salt and minerals, frequently in association with iron compound. Magnesium is vital micronutrient for both plant and animal. Magnesium is often associated with calcium in all kind of water, but its concentration remain generally lower than the calcium [35]. Magnesium is essential for chlorophyll growth and act as a limiting factor for the growth of phytoplankton [18]. In Chandlodia lake the amount of magnesium recorded ranges between 30 ± 2.58 to 34 ± 2.12 . The maximum amount of magnesium in the water was recorded during winter season and minimum amount was recorded during summer season

Decrease in level of magnesium reduces the phytoplankton population. [36] suggested that the considerable amount of magnesium influence water quality. Magnesium is essential for chlorophyll bearing plant. Magnesium enters into combination with anions other than CO_2 in lakes such as chloride

and sulphate [37]. Various sub-processes like bating, picking, tanning, dyeing and fat liquoring causes water pollution [38]. Magnesium is vital micronutrient for both plant and animal.

3.10 Dissolved Oxygen

The oxygen in water can be dissolved from air or is produced from the photosynthetic organism like algae and aquatic plants oxygen is poorly soluble gas in water and its solubility depend on the temperature of water and its partial pressure. The solubility of oxygen also decreases with increasing salinity of water. [39] has established a direct relationship between photosynthesis and dissolved oxygen. Measurement of dissolved oxygen is a primary parameter in all pollution studies. The amount of dissolved oxygen recorded in the water of Chandlodia lake ranges between 2.14 ± 0.07 to 5.08 ± 0.07 . The minimum amount of dissolved oxygen in the water of Chandlodia lake was recorded during winter season whereas maximum amount was recorded during monsoon season

The high temperature and addition of sewage and other waste might be responsible for low value of DO [40] [41]. Depletion of dissolved oxygen in water is due to high temperature and increased microbial activity [42]. Dissolved oxygen with high value observed during monsoon may be as a result of the increased solubility of oxygen at lower temperature [43].

3.11 Biochemical oxygen demand

BOD refers the oxygen used by the microorganism in the aerobic oxidation of organic matter. Therefore with the increase in the amount of organic matter in the water the BOD increases. The BOD value in Chandlodia lake ranges between 1.96 ± 0.08 to 3.38 ± 0.3 . The minimum demand of oxygen in the water was recorded during summer season, whereas maximum demand was recorded during monsoon season

The higher value of BOD during monsoon was due to input of organic wastes and enhanced bacterial activity [44]. The reason of high BOD in monsoon might also be due to presence of several microbes in water bodies, which accelerate their metabolic activities with the increase in concentration of organic matter in the form of municipal and domestic waste which was discharge into water bodies and so the demand of oxygen increased [45].

3.12 Chloride

The greater source of chlorides in lake water is disposal of sewage and industrial waste. Human body release very high quantity of chlorides through urine and faeces. The chloride concentration was used as an important parameter for detection of contamination by sewage. Prior to development of bacteriological and other test like BOD and COD. The amount of chloride recorded in the water of

Chandlodia lake ranges between 88 ± 4.97 to 122 ± 4.97 . The minimum amount of chloride in the lake water was recorded during winter season and the maximum amount was recorded during summer season.

The higher concentration of chloride during summer month may be associated with frequently run-off loaded with contaminated water from the surrounding [46] [47]. The high chloride concentration of the lake water may be due to high rate of evaporation [48] or due to organic waste of animal origin [49].

3.13 Sodium

Sodium is a natural constituent of raw water, but its concentration is increased by pollutional sources such as rock salt, precipitation runoff, soapy solution and detergent. The amount of sodium recorded in the water of Chandlodia lake ranges between 38 ± 4.97 to 65 ± 2.89 . The minimum amount of sodium in the water of Chandlodia lake was recorded during winter and maximum amount was recorded during monsoon season

The high level of sodium during monsoon may be attributed to the rain water as it carries the salt dissolved from the surrounding area [50]. The addition of waste water containing soap solution and detergent from the surrounding slummy area are also responsible for the increase in sodium level in the water bodies.

3.14 Nitrate

Nitrates are contributed to freshwater through discharge of sewage and industrial wastes and run off from agricultural fields. Some ground waters naturally have high nitrate concentration. In Chandlodia lake the amount of nitrate recorded ranges between 5.8 ± 0.18 to 7.4 ± 0.22 . The minimum amount of nitrate in the water of Chandlodia lake was recorded during summer season, whereas maximum amount of nitrate in the water was recorded during monsoon season.

The high nitrate concentration during monsoon might be due to influx nitrogen rich flood water that brings about large amount of contaminated sewage waste. The monsoon season was the period with the highest nitrate-nitrogen concentration which is known to support the formation of blooms [51] [52] [53]. Nitrate content was higher in monsoon season, which can be attributed to the nitrate leached from the surrounding area. But lower concentration in summer was due to utilization by plankton and aquatic plants [54].

3.15 Phosphate

Excess amount of phosphate may cause eutrophication leading to extensive algal growth called algal blooms. Total phosphates in water include both organic and inorganic phosphates. Organic phosphates are part of living and dead plants

and animal; over 85% of total phosphorous is usually found in organic form. In Chandlodia lake the amount of phosphate recorded ranges between 1.24 ± 0.05 to 1.92 ± 0.11 . The minimum amount of phosphate recorded in the water of lake was during winter season and the maximum amount was recorded during monsoon season.

The maximum value of phosphate during monsoon may be attributed to surface runoff during rainy season receiving huge quantity of domestic sewage, cattle dung and detergents from the surrounding catchment area. Catchment area activities are enriching phosphate in the lake [55]. The lower value of phosphate in summer month may be due to more uptake of phosphate for luxuriant growth of macrophytes.

4. CONCLUSION

Therefore comparing the result with [56] [57] standard it was concluded that water of Chandlodia lake shows very huge level of pollution and similar condition continue the lake with soon become ecological inactive.

REFERENCES

- [1] APHA, AWWA. *Standard. Methods for the examination of water & waste water.* (Washington DC 18th Edition, 1985).
- [2] S.M. Kumar and S. Ravindranath, "Water Studies – Methods for monitoring water quality". (Banglore, Karnataka, Center for Environment Education (CEE), 1998).
- [3] R.K.Trivedy and P.K. Goel, *Chemical and biological methods for water pollution studies.* (Karad, Maharashtra (India), Environmental Publication,1984).
- [4] B.K. Dwivedi and G.C. Pandey, Physico-chemical factors and algal diversity of two ponds in Faizabad, India *Poll.Res.* 21(3), 2002, 361-370.
- [5] R.P. Singh and P. Mathur, Investigation of variations in physicochemical characteristics of a fresh water reservoir of Ajmer city, Rajasthan, *Ind. J. Environ. Science*, 9, 2005, 57-61.
- [6] P.S. Welch, *Limnology*, (N.Y. 2nd Ed., McGraw Hill Book Co.,1952).
- [7] P.C. Joshi and A. Singh, Analysis of certain physicochemical parameters and plankton of freshwater hill stream at Nanda devi biosphere reserve. *Uttarpradesh J. Zool.*, 21, 2001, 177-179.
- [8] B.B. Ghose and A.K. Basu, Observation on estuarine pollution of Hoogly by the effluents from a chemical factory complex at Reshase, West Bengal. *Journal of Env. Health.* 10, 1968, 209-218.
- [9] J.O. Young, Seasonal and diurnal changes in the water temperature of a temperate pond

- (England) and tropical pond (Kenya). *Hydrobiol.* 47, 1975, 513-526
- [10] H. S. Sehgal, *Limnology of lake Sruinsar, Jammu with reference to zooplankton and fisheries prospectus*, doctoral diss., University of Jammu, 1980.
- [11] M. Jayanti, *A comprehensive study of three contrasting lentic system in the content of Aquaculture*, doctoral diss., Bharathidasan University, Tiruchirappalli, 1994.
- [12] A.A. Ahluwalia, *Limnological Study of wetlands under Sardar Sarovar command area*. doctoral diss., Gujarat University, Ahmedabad, 1999.
- [13] H.A. Solanki and B.R. Pandit, Trophic status of lentic waters of ponds water of Vadodara, Gujarat state, India. *Int. J. of Bioscience Reporter* 4 (1), 2006, 191-198.
- [14] A.K.Das and N.P. Shrivastva, Ecology of Sarni Reservoir (M. P.) in the context of Fisheries. *Poll Res.* 22(4), 2003, 533-539.
- [15] K.L. Saxena, R.N. Chakraborty, A.Q. Khan and S.N. Chattopadhyaya, Pollution studies of the river Ganga near Kanpur. *Indian J. Environ. Hlth.* 8, 1966, 270-285.
- [16] K.K. Ansari, and S. Prakash, Limnological studies on Tulsidas Tal of Tarai region of Balrampur in relation to fisheries *Poll. Res.* 19(4), 2000, 651-655.
- [17] H.A. Solanki, *Study on pollution of soils and water reservoirs near industrial areas of Baroda*, doctoral diss., submitted to Bhavnagar University, Bhavnagar, 2001.
- [18] Dagaonkar and D.N. Saksena, Physicochemical and biological characterization of a temple tank, Kaila Sagar, Gwalior, Madhya Pradesh. *J. Hydrobiol.* 8 (1), 1992, 11-19.
- [19] R.K. Garg, R.J. Rao and D.N. Saksena, Assesment of physicochemical water quality of Harsi Reservoir, District Gwalior, Madhya Pradesh. *J. Ecophysiol. Occupat. Health* 6, 2006, 33-40.
- [20] N. Swarnalatha and A. Narasingrao, Ecological studies of Banjara lake with reference to water pollution. *J. Environ. biol.* 19 (2), 1998, 179-186.
- [21] K.D. Sharma, N. Lal and P.D. Pathak, Water quality of sewage drains entering Yamuna at Balasurami, Physicochemical and Bacteriological Analysis of water from Namakkal and Erode Districts, Tamilnadu, India. *Poll Res.* 25(3), 2006, 495-498.
- [35] R. Venkatasubramani and T. Meenambal, Study of subsurface water quality in Mattupalayam Taluk of Coimbatore district Tamil Nadu. *Nat. Environ. Poll. Tech.* 6, 2007, 307-310
- [36] V.S. Govindan and R. Devika, Studies on Heavy metal profiles of Adyar river and Agra. *Indian J. Environ Hlth*, 23, 1981, 118-122.
- [22] M.G. George, Diurnal variation in two shallow ponds in Delhi , India. *Hydrobiol.*, 3, 1962, 265.
- [23] A.M. McCombie, Factors Influencing the growth of phytoplankton Canada. *J. Fish Res. Ed.*, 10, 1953, 253-282.
- [24] S.N. Nandan and R.J. Patel, *Ecological studies of algae in aquatic ecology* (New Delhi, Ashis Publishing House, 1992).
- [25] S.K. Moitra and B.K. Bhattacharya, Some hydrological factors affecting plankton production in fish pond in Kalyani, West Bengal, India. *Ichthyologia* 4 (1 & 2), 1965, 8-12.
- [26] I.A. Wani and Subla, Physicochemical features of two shallow Himalayan lakes. *Bull Eviron. Sci.*, 8, 1990, 33-49.
- [27] M. Chaurasia and G.C. Pandey, Study of physico- chemical characteristic of some water pond of Ayodhya –Faizabad. *Indian J. of Environmental protection*, 27 (11), 2007, 1019-1023.
- [28] S.A. Abbasi, F.J. Khan, K. Sentilevelan and A. Shabuden, *Indian J. Env. Hlth.* 14(3), 1999, 176-183
- [29] C.K. Jain, K.K.S. Bhatia and T. Vijay, Ground water quality in coastal region of Andhra Pradesh. *Indian J. of Environmental Health.* 39 (3), 1997,182-190.
- [30] S. Padma and Periakali, Physicochemical and geochemical studies in Pulicat lake, east coast of India. *Indian J.Mar. Sci.*, 28, 1999, 434-437.
- [31] N.K. Patel and B.K. Sinha, Study of the pollution load in the pond of Burla area near Hirakund dam at Orissa. *J. Env. Poll.* 5, 1998, 157-160.
- [32] S.B. Angadi, N. Shiddamaliayya and P.C. Patil, Limnological study of papnash pond, Bidar (Karnataka). *J. Env. Biol.*, 26, 2005, 213-216.
- [33] A.K. Tripathi and S.N. Pandey, *Water pollution* (New Delhi, Ashish Publishing House, 1990).
- [34] J. UdhayaKumar, D. Natarajan, K. Srinivasan, C. Mohansundari, and M. waste stabilization pond. *J. Ecotoxicol. Environ. Monit.*, 1(1), 1991,53-58.
- [38] V.G. Jhingran, *Fish and Fisheries of India* (Delhi (India), Hindustan Publ. Corp., 1975).
- [39] R.L. Bolton and L. Klein, *Sewage treatment basic principles and trends* (London, Butterworths, 1971).
- [40] S. Vijayaraghavan, Seasonal variation in pri. Productivity in three tropical ponds. *Hydrobiol.* 38, 1971, 395-408.


- [41] G.M. Woodward, Pollution control in Hamber estuary. *Water pollution control* . 83 (1), 1984, 82-90.
- [42] A.S. Mathuthu, F.M. ZaranYika, S.B. Jannalagadde, Monitoring of water quality in upper Mukavisi river in Harare, Zimbabwe. *Environ . int.* 19(1), 1993, 51-56.
- [43] H.C. Kataria, A. Singh and S.C. Pandey, Studies on water quality of Dahod Dam, India. *Poll. Res.* 25(3), 2006, 553–556.
- [44] A.A. Prasannakumari, T. Ganagadevi and C.P. Sukeshkumar, Surface water quality of river Neyyar- Thiruvananthapuram, Kerala, India. *Poll Res.* 22(4), 2003, 515 – 525.
- [45] S. Kaushik and D.N. Saksena, Physico chemical limnology of certain waterbodies of central India. in. Kvismayan (Ed.), *Freshwater ecosystem in India*. (Delhi: Daya Publishing House, 1999) 336.
- [46] June and C. Fred, Physical, chemical and biological characteristics of lake Sharpe, South Dakota (USA). *US Fish wild. Serv. Fish wild L., Tech. rep.* 0(8), 1987, 1-20.
- [47] S. Sunder, Mounting the water quality in a stretch of river Jhelum. Kashmir, in Book” *Ecol and Poll. Of Indian rivers* (New Delhi: Ashish Publishing House, 1988) 131-161
- [48] A. Kumar, Observation on the diel variations in abiotic and biotic components of the river Mayurrakshi (Santal Pargana). Bihar. *Indian .J. Ecol.* 22 (1), 1995, 39-43.
- [49] B.N. Prasad, Y.C. Jaitly and Y. Singh, Periodicity and interrelationships of physico-chemical factors in ponds. In: A.D. Adoni (eds.) *Proc. Nat. Symp. Pure and Apply. Gmnol. Bull. Bot. Soc. Sagar.*, 32, 1985, 1-11.
- [50] S.S. Purohit, and M.M. Saxena, *In water, Life and Pollution.*(Bikana, ed.): India, Arg. Botanical Publishers 1990).
- [51] R. Sahai and A.B. Sinha, Investigation on bio-ecology of inland water of Gorakhpur (UP), India. I. Limnology of Ramgarh Lake. *Hydrobiol.* 34(3), 1969, 143-447.
- [52] P. Blomqvist, A. Petterson and Hyenstrand, Ammonium –nitrogen: A key regulatory factor causing dominance of non nitrogen fixing cyanobacteria in aquatic systems. *Arch. Hydrobiol.*, 132, 1994, 141-164.
- [53] D.M. Anderson, A.D. Cembella and G.M. Hallegraeff, *Physiological Ecology of Harmful algal blooms. 1st Edn.*, (Berlin: Springer-Verlag, 1998) 647-648.
- [54] P.V. Zimba, L. Khoo, P.S. Gaunt, S. Brittain, and Carmichael, Confirmation of cat fish, *Ictalurus punctatus* Ralfinesque, mortality from microcystis toxins. *J. Fish Dis.*, 24, 2001, 41-47.
- [55] V. Kannan, *The limnology of Sathiar: A freshwater impoundment*, doctoral diss., Madurai Kamraj University, Madurai, 1978.
- [56] S. Tomat and P. Sharma, Physico chemical status of Upper lake (Bhopal, India). Water quality with special reference to phosphate and nitrate concentration and their impact on lake ecosystem. *Asian .J. Exp.Sci.* 20 (2), 2006, 289-296.
- [57] WHO, *International Standards for drinking water*, (3rd Ed. Geneva, World Health Organization, 1971).
- [58] BIS, *Standard tolerance limits for bathing water* (India, Bureau of Indian Standards, 1982).

RESULT TABLE: CHANDLODIA LAKE


Sr. no.	PARAMETERS	2010 - 2011		
		Summer	Monsoon	Winter
		Mean + S.E.	Mean + S.E.	Mean + S.E.
1.	Temperature in °C	29 ±1.58	21 ±1.29	16 ±1.47
2.	Electrical conductivity in mhos/cm	2.78 ±0.07	3.24 ±0.26	2.94 ±0.07
3.	Turbidity in NTU	16 ±1.47	22 ±0.91	18 ±1.58
4.	Total Dissolve Solid in ppm	1014 ±37	1264 ±39	963 ±31.5
5.	pH	8.7 ±0.17	9.1 ±0.18	8.9 ±0.15
6.	Alkalinity in ppm	192 ±3.65	214 ±6.87	208 ±5.6
7.	Total Hardness in ppm	312 ±5.89	328 ±8.29	304 ±6.58

8.	Calcium in ppm	76 ±3.92	82 ±4.32	66 ±3.92
9.	Magnesium in ppm	30 ±2.58	31 ±1.73	34 ±2.12
10.	Dissolved Oxygen in ppm	2.14 ±0.07	5.08 ±0.07	3.28 ±0.22
11.	Biochemical Oxygen Demand in ppm	1.96 ±0.08	3.38 ±0.3	2.20 ±0.21
12.	Chloride in ppm	122 ±4.97	93 ±4.65	88 ±4.97
13.	Sodium in ppm	44 ±3.65	65 ±2.89	38 ±4.97
14.	Nitrate in ppm	5.8 ±0.18	7.4 ±0.22	6.7 ±0.32
15.	Phosphate in ppm	1.66 ±0.12	1.92 ±0.11	1.24 ±0.05

[S.E. = Standard error]



SATELLITE IMAGE



REAL IMAGE